



**US Army Corps
of Engineers**

Hydrologic Engineering Center

HEC-EFM

Ecosystem Functions Model

Quick Start Guide

Version 2.0
November 2009

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to the Department of Defense, Executive Services and Communications Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.

1. REPORT DATE (DD-MM-YYYY) November 2009			2. REPORT TYPE Computer Program Documentation		3. DATES COVERED (From - To)		
4. TITLE AND SUBTITLE HEC-EFM Ecosystem Functions Model Quick Start Guide Version 2.0				5a. CONTRACT NUMBER			
				5b. GRANT NUMBER			
				5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S) John Hickey Woody Fields				5d. PROJECT NUMBER			
				5e. TASK NUMBER			
				5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Corps of Engineers Institute for Water Resources Hydrologic Engineering Center (HEC) 609 Second Street Davis, CA 95616-4687				8. PERFORMING ORGANIZATION REPORT NUMBER CPD-80a			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/ MONITOR'S ACRONYM(S)			
				11. SPONSOR/ MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.							
13. SUPPLEMENTARY NOTES Also, see HEC-EFM User's Manual, CPD-80							
14. ABSTRACT The Ecosystem Functions Model (HEC-EFM) is a planning tool that aids in analyzing ecosystem response to changes in flow regime. The Hydrologic Engineering Center (HEC) of the U.S. Army Corps of Engineers is developing HEC-EFM to enable project teams to visualize existing ecologic conditions, highlight promising restoration sites, and assess and rank alternatives according to the relative change in ecosystem aspects.							
15. SUBJECT TERMS HEC-EFM, Ecosystem Functions Model, ecosystem modeling, ecosystem restoration, time series analysis, hydraulic modeling, Geographic Information Systems, river and wetland flow regimes, ecological relationships							
16. SECURITY CLASSIFICATION OF:				17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT	b. ABSTRACT	c. THIS PAGE					19b. TELEPHONE NUMBER
Unclassified	Unclassified	Unclassified	Unlimited		48		

HEC-EFM

Ecosystem Functions Model

Quick Start Guide

Version 2.0
November 2009

US Army Corps of Engineers
Institute for Water Resources
Hydrologic Engineering Center
609 Second Street
Davis, CA 95616

(530) 756-1104
(530) 756-8250 FAX
www.hec.usace.army.mil

CPD-80a

Ecosystem Functions Model, HEC-EFM, Quick Start Guide

2009. This Hydrologic Engineering Center (HEC) documentation was developed with U.S. Federal Government resources and is therefore in the public domain. It may be used, copied, distributed, or redistributed freely. However, it is requested that HEC be given appropriate acknowledgment in any subsequent use of this work.

Use of the software described by this document is controlled by certain terms and conditions. The user must acknowledge and agree to be bound by the terms and conditions of usage before the software can be installed or used. For reference, a copy of the terms and conditions of usage are included in Chapter 5 of this document so that they may be examined before obtaining and loading the software. The software described by this document can be downloaded for free from our internet site (www.hec.usace.army.mil).

HEC cannot provide technical support for this software to non-Corps users. See our software vendor list (on our web page) to locate organizations that provide the program, documentation, and support services for a fee. However, we will respond to all documented instances of program errors. Documented errors are bugs in the software due to programming mistakes not model problems due to user-entered data.

This document contains references to product names that are trademarks or registered trademarks of their respective owners. Use of specific product names does not imply official or unofficial endorsement. Product names are used solely for the purpose of identifying products available in the public market place.

Microsoft and *Windows* are registered trademarks of Microsoft Corp.

ArcGIS, *ArcView*, and *ArcInfo* are registered trademarks of ESRI, Inc.

Table of Contents

Chapters

1	Introduction.....	1-1
2	User Interface.....	2-1
2.1	Properties	2-2
2.2	Relationships.....	2-3
2.3	Tables.....	2-5
2.4	Combo Relationships.....	2-6
3	HEC-EFM Math	3-1
3.1	Season	3-1
3.2	Duration	3-1
3.3	Rate of Change.....	3-2
3.4	Time Series Specifications.....	3-3
4	Demonstration Project.....	4-1
4.1	Building Relationships.....	4-1
4.1.1	Little minnow spawning habitat.....	4-1
4.1.2	Big bass winter habitat.....	4-2
4.1.3	Benthic macro invertebrate biodiversity.....	4-2
4.1.4	Wetland health	4-3
4.1.5	Riparian tree recruitment and inundation	4-3
4.2	Setting up the Model.....	4-5
4.2.1	Creating a New Project and Defining Flow Regimes.....	4-5
4.2.2	Defining Relationships.....	4-8
4.2.3	Results.....	4-12
4.2.4	HEC-EFM Output.....	4-15
4.2.5	Using HEC-EFM Plotter.....	4-18
4.3	Hydraulic Analysis.....	4-20
4.3.1	HEC-RAS	4-21
4.3.2	HEC-GeoRAS.....	4-22
4.4	HEC-EFM Analyses with GIS.....	4-22
5	Conclusions.....	5-1
6	Terms and Conditions for Use.....	6-1

CHAPTER 1

Introduction

The Ecosystem Functions Model (HEC-EFM) is a planning tool that aids in analyzing ecosystem response to changes in flow regime. The Hydrologic Engineering Center (HEC) of the U.S. Army Corps of Engineers is developing HEC-EFM to enable project teams to visualize existing ecologic conditions, highlight promising restoration sites, and assess and rank alternatives according to the relative change in ecosystem aspects.

Central to HEC-EFM analyses are "functional relationships". These relationships link characteristics of hydrologic and hydraulic time series (flow and stage) to elements of the ecosystem through combination of four basic criteria: 1) season, 2) duration, 3) rate of change, and 4) percent exceedance.

After relationships are developed, HEC-EFM performs statistical computations to analyze flow and stage time series for the specified criteria and produces a single flow value for each relationship. This process can then be repeated for alternative flow regimes to compare different project scenarios and indicate directions of changes to ecosystem health.

In addition to the statistical computations, HEC-EFM analyses typically involves hydraulic modeling, which can translate statistical results to water surface profiles and spatial layers of water depth, velocity, and inundation areas. Geographic Information Systems (GIS) can then be used to display these generated layers as well as other relevant spatial data (i.e., soils, vegetation, and land-use maps).

Data requirements of HEC-EFM are related to the level of detail desired by the modeler. If only statistical results are desired, then required data consists of the flow regimes to be analyzed and the eco-hydro relationships. If the user intends to visualize statistical results spatially, data (and software) requirements increase significantly to include flow and stage time series, eco-hydro relationships, digital topography, a geo-referenced hydraulic model, and any other spatial data relevant to the ecosystem investigations.

This quick start guide provides an overview of the software followed by examples that demonstrate its application. The text has been formatted to help readers keep track of the different types of information presented. *Italics* are used to identify *software features* that are available through the user interfaces of HEC-EFM. Underlines are used to identify model input data, which includes the names of flow regimes and relationships used in the demonstration project. **Bold** is used to highlight **key information** for individual sections of text.

CHAPTER 2

User Interface

The main interface of HEC-EFM consists of a title bar, a series of menus, four tabs (*Properties*, *Relationships*, *Tables*, and *Combo Relationships*), a message bar, and a *Recalculate* button (Figure 1). Most features are located on the tabs. The rest of this section provides more detail for those tabs.




Figure 1. Key components of the main interface of HEC-EFM.

2.1 Properties

The *Properties Tab* contains information relevant to the model as a whole and is divided into two frames: *Model Information* and *Flow Regimes* (Figure 2).




Figure 2. The *Properties Tab* of the main interface of HEC-EFM is used to document information about the project and to define the flow regimes to be analyzed. Two flow regimes (**Natural** and **Gaged**) are shown.

Model Information – *Model information* documents metadata (*Title, Author, Location, and Description*) about the project.

Flow Regimes – An HEC-EFM "flow regime" is defined as two concurrent daily time series that reflect conditions at a single location in the study area. Typically, a flow regime is composed of time series of daily mean flow and daily mean stage data. HEC-EFM accepts input data from several text formats (i.e., comma, space, and tab delimited) and from HEC Data Storage System

(HEC-DSS), which is the database used by HEC models for storage of time series and other data. The start and end dates for flow regimes are controlled by the user, which can be a handy feature when analyzing only part of the period of record. **HEC-EFM does not allow missing data.**

2.2 Relationships

HEC-EFM "relationships" are statistical representations of links between hydrology and ecology. Relationships are typically developed by teams of scientists and engineers using a combination of expert knowledge and scientific literature. Relationships are defined on the *Relationships Tab*, which is broken up into four frames: *Statistical queries*, *Geographical queries*, *Other queries (nonstandard)*, and *Options* (Figure 3).




Figure 3. Using the *Relationships Tab* of the main interface of HEC-EFM, modelers define individual relationships as combinations of statistical and geographical criteria.

Statistical Queries – Fields within *Statistical queries* allow users to input criteria (in terms of *Season*, *Duration*, *Rate of change*, and *Percent exceedance*) that define the statistical analysis to be performed for each relationship. All criteria do not need to be filled in for each relationship – only those that are important for the relationship. Time series specifications can be entered as *Flow frequency* (percent of years) or *Flow duration* (percent of time). *Statistical queries* also offer controls for managing the flow and stage data to be used for the statistical computations. Options are given for *Water year range*, *Individual water year*, or *Relationship-defined water year*, which allows a relationship to be based on the statistical results of a separate relationship.

Geographical Queries – *Geographical queries* allow users to specify criteria that define relationships from a spatial perspective. Geographical queries are created by the user through the *Manage Tags for Geographical Queries* interface (Figure 4), which is accessed by selecting the "Edit – Manage Geo Queries..." menu option. Users create tags and specify the tags that are important for each relationship. This creates a text field in the HEC-EFM interface for recording how the tagged data sets need to be queried in GIS. A tag is a simple name that identifies sets of spatial data layers important to an HEC-EFM project.




Figure 4. The *Manage Tags for Geographical Queries* interface in HEC-EFM is used for editing and creating tags, which are labels for sets of spatial data important to an HEC-EFM project.

Other Queries (nonstandard) – Statistical results for a standard HEC-EFM relationship is a pair of flow and stage data that meet the statistical criteria specified for that relationship. Currently, the only nonstandard query available is the *reverse lookup query*. Reverse lookups do not compute a flow and stage, instead the user specifies a flow and HEC-EFM computes the percent of years or percent of time that flow is equaled or exceeded.

Options – There are three options available for each relationship: *Write computation arrays*, *Hypothesis tracking*, and *Confidence tracking*. The *Write computation arrays* option prompts HEC-EFM to export the statistical computations performed for that relationship. *Hypothesis tracking* allows users to compare the direction of change of eco-health for different flow regimes. The question that hypothesis tracking asks is: "Increased flow will do what to eco-health?" for this relationship. Users have the option of saying that increased flow will help (+), hurt (-), or have a non-linear response (via the *Curve* button) to eco-health. *Confidence tracking* provides a way to track the relative certainty of HEC-EFM relationships. The default is one star. This starting point implies that there is the same amount of scientific understanding for each of the relationships. As confidence in a particular relationship grows, its number of stars can be increased. This occurs at the discretion of the study team, perhaps when the relationship is verified with field data, backed with scientific literature, or approved by a group of scientists or agencies. A common result when analyzing multiple relationships is that for a certain flow regime, some are helped and some are hurt. *Indices* are a tool that can help users look at the net effect of different flow regimes. *Confidence tracking* and *hypothesis tracking* must be in use if you want to include a particular relationship in an index. Indices are computed for flow regimes that are active and not the reference. Each index is computed using a combination of information about relationships and statistical results based on the following equation:

$$\text{Index} = \sum_{i=1..n} (\text{Direction of Change}_i) * (\text{Confidence}_i) * (\% \text{ Change in Eco-value}_i)$$

Where:

i = counter from 1 to n

n = number of relationships in the index

Direction of Change_i = 1, -1, or 0 for relationship_i

1 indicates that relationship_i experienced a positive change from the reference flow regime

-1 indicates that relationship_i experienced a negative change from the reference flow regime

0 indicates that relationship_i experienced no change from the reference flow regime

Confidence_i = an integer from 0 to 5 based on the confidence value for relationship_i

% Change in Eco-value_i = $\frac{(\text{Eco-value for relationship}_i - \text{Eco-value for relationship}_{\text{reference}})}{(\text{Eco-value for relationship}_{\text{reference}})}$

Note: % change in Eco-value is equal to % change in Flow (using statistical results for flow) when hypothesis tracking does not use the curve option.

2.3 Tables

The *Tables Tab* presents statistical results for pairings of flow regimes and relationships. It is also where users can view project reports and output files (Figure 5). Tables can be printed or copied into other applications. After the *Recalculate* button is pressed, the HEC-EFM interface will automatically go to the *Tables Tab* to display the new statistical results.





Figure 5. The *Tables Tab* of the HEC-EFM interface is used to display statistical results for simulations (as in this figure), project reports, and computational output.

2.4 Combo Relationships

Combo Relationships are used to detail how two or more individual relationships are grouped to represent a single ecosystem dynamic. For example, in Figure 6, Riparian tree establishment is defined as Riparian tree recruitment minus Riparian tree inundation.

Figure 6. The *Combo Relationships Tab* of the HEC-EFM interface allows for grouping of separate relationships to represent a single ecosystem dynamic.



CHAPTER 3

HEC-EFM Math

While the HEC-EFM process can involve hydraulic modeling and GIS, statistical analyses of links between hydrology and ecology are the foundation of HEC-EFM applications.

The math performed by HEC-EFM during these statistical analyses is actually very basic. The challenge is in understanding how the parameters that define relationships (*Season*, *Duration*, *Rate of change*, and *Percent exceedance*) are used by HEC-EFM to compute the flow and stage that meet those criteria.

3.1 Season

Season is the most intuitive of the parameters. Ecosystem dynamics (e.g., fish spawning or seed germination) typically occur in certain time periods of the year. Within HEC-EFM, season is defined by start and end dates. During computations, a seasonal extract is taken from each water year being investigated. All *Duration* and *Rate of change* queries are performed on these seasonal extracts.

3.2 Duration

Duration is a versatile, but complicated query. There are three settings: 1) *number of days*, 2) a selection of statistics to be computed *for each duration* in the season, and 3) a selection of statistics to be computed using the time series of *computed values* produced in step two.

The *number of days* defines a duration interval (Figure 7). *Duration* calculations are performed from the beginning of season to the end of season. So for the start date, HEC-EFM considers all data values within the duration interval (per setting 1), computes a statistic (per setting 2), records that value for the start date, and then advances a day and repeats the process until the end of season is reached. This produces a statistical time series of *Minimums*, *Medians*, *Maximums*, *User defined percentages*, or *Means* (per setting 2) that has one value for each day of the season.

The final step in the *Duration* query involves selecting a single seasonal value (per setting 3) from the statistical time series. Options are provided for *Minimum*, *Median*, *Maximum*, *User defined percentage*, and *Mean*. This produces a time series of seasonal results that has one value per season.

Figure 7 provides an example of this process using an 8 day duration interval, minimums, and then maximum of the minimums.




Figure 7. Sample use of the *Duration* query in HEC-EFM. Computations are shown for one water year.

3.3 Rate of Change

The *Rate of change* query was originally added to HEC-EFM to look at ecological concerns connected to the recession limb of stage hydrographs. Its capabilities have since been expanded to include *Rising*, *Falling*, and *Absolute* rates of change for both *Stage* and *Flow*.

For all combinations of those settings (Figure 8), the *Rate of change* uses two parameters: 1) the *threshold value for change* and 2) the *number of days*. For each day beginning at the end of the season and working backwards in time, the actual rate of change is computed by subtracting the current flow or stage and the flow or stage that occurs at the end of the time interval defined by the *number of days* (i.e., if *number of days* is set to 4, the actual rate of change equals the current value minus the value 3 days into the future such that the total time interval being considered spans 4 days). The actual rate of change is then compared to the *threshold value*. If actual does not violate the *threshold*, the rate of change is deemed acceptable, HEC-EFM moves backwards one day, and the test is repeated. This continues until the *threshold* is violated or the beginning of season is reached.

If the *threshold* is violated, HEC-EFM selects the previous successful test (one day later than the failed test) as the seasonal result, which represents the date and conditions where rates of change became consistently acceptable for the rest of the season.

If the beginning of season is reached and passes the rate of change test, HEC-EFM selects that date and its corresponding conditions as the seasonal result.




Figure 8. Sample use of the *Rate of change* query in HEC-EFM. Computations are shown for one water year.

If the first test (at the end of season) fails, a zero value is recorded as the seasonal result, which indicates that no dates passed the rate of change test for that season.

It is also important to note that when *Rate of change* is selected in the HEC-EFM interface, the dropdown controls in the *Duration* query become gray and inoperable. This behavior is meant to communicate that the two queries are largely incompatible. In fact, the only *Duration* setting relevant to *Rate of change* is *number of days*. When it is set to 1 day (as in Figure 8), the *Rate of change* query tests the mean daily values obtained from the flow regimes in the seasonal extract. When an integer of 2 or more is entered, HEC-EFM will compute a time series of mean values for that duration of flow or stage, depending on the setting in *Rate of change*, and then perform the rate of change tests for that computed series.

3.4 Time Series Specifications

Components in the *Time series specifications* section of HEC-EFM include *Percent exceedance* and three controls for managing the period of record assessed for each relationship (*Water year range*, *Individual water year*, and *Relationship-defined water year*).

The *Percent exceedance* query offers a choice of either *Flow frequency* or *Flow duration*. *Flow duration* should not be confused with the *Duration* query described earlier in this chapter. In fact, when *Flow duration* is selected in the HEC-EFM interface, the dropdown controls in the *Duration* query become gray and inoperable. As with *Rate of change*, the only *Duration* setting relevant to *Flow duration* is *Number of days*. When it is set to 1 day, HEC-EFM generates a flow duration curve using mean daily values obtained from the flow regimes in the seasonal

extract and then interpolates to obtain the flow that corresponds to the user-defined percentage. When an integer of 2 or more is entered, HEC-EFM computes a time series of mean flow values for that duration, uses those values to generate a flow duration curve, and then interpolates to obtain the flow that is equaled or exceeded for the **user-defined percentage of time**. The resulting value would be the statistical result.

When *Flow frequency* is selected (Figure 9), HEC-EFM ranks the seasonal results (computed via the *Season*, *Duration*, and *Rate of change* queries) and interpolates to obtain the flow (or stage, if *Rate of change* is being used to investigate stage dynamics) that is equaled or exceeded for the **user-defined percentage of years**. The resulting value would be the statistical result.




Figure 9. Sample use of the *Time series specifications* for a relationship with *Season* and *Duration* queries (as in Figure 7). The statistical result is the flow meeting the parameters in those queries that is equaled or exceeded in 25% of years (*Flow frequency*).

This ranking and interpolation process is required only when seasonal results are computed for multiple years, whether for a whole flow regime as defined on the Properties tab (Figure 2) or via the *Water year range* control (Figure 9).

Both the *Individual water year* and the *Relationship-defined water year* controls limit analyses to a single water year. For the *Individual water year*, the user specifies which water year to assess. For *Relationship-defined water year*, the user selects a separate relationship that is of interest, and HEC-EFM determines which of its seasonal results (and corresponding historical water year) most closely equals its statistical result. That historical water year is then used when assessing the current relationship. In both cases, as there is only one seasonal result, it is also reported as the statistical result.

CHAPTER 4

Demonstration Project

This demonstration project will consider ecosystem relationships for a cross section of the fictional Rolling River. The progression of tasks is to 1) build eco-hydro relationships, 2) set up the model and compute statistical results, 3) simulate the statistical results with a hydraulic model, and 4) use of GIS to investigate the ecosystem responses spatially.

Two flow regimes will be analyzed. The "Gaged" flow regime has flows and stages that reflect the current management plan for Rolling River, which is regulated by upstream dams. The "Natural" flow regime has flows and stages that reflect how the river would behave without reservoir regulation.

A completed demonstration project with flow regime data can be downloaded via the "*Help – Install Demonstration Project...*" menu option.

4.1 Building Relationships

Each scenario described below presents information about an aspect of the ecosystem that the Rolling River study team would like to investigate, the statistical and geographical queries used to define each as an HEC-EFM relationship, and the logic used to craft those criteria.

4.1.1 Little minnow spawning habitat

The little minnow is a threatened species of fish indigenous only to the Rolling River basin. Little minnow populations have declined (in magnitude and range) as dams and diversions limited access to upstream river stretches and as floodplain areas, critical to little minnow spawning, were transitioned to agricultural lands. Little minnow spawn in **shallow (no deeper than 3 ft), vegetated** floodplain areas between **February and May**. Eggs require **sustained high flows** for approximately **21 to 28 days** before hatching. Little minnow reach sexual maturity in their first or second year and have a lifespan of approximately 6 years. Scientists suggest using the conditions within each spawning season that generate the **largest extent** of effective spawning habitat as an indicator of success for each year's spawn. Further, scientists suggest that good spawning conditions do not need to occur every year – it would be sufficient if there were good conditions in **25% of years**, so that, on average, each little minnow would have a chance to spawn in their lifespan.

HEC-EFM Relationship:

- *Season:* 02/01 to 05/31
- *Duration:* 24 days, *Minimums* (sustained highs) and then *Maximum* (largest extent)
- *Rate of change:* Not applied
- *Percent exceedance:* 25% (4 yr) - *Flow frequency*

- *Hypothesis tracking*: Increased flow will improve (+) floodplain spawning
- *Geographical queries*: Depth (0 to 3 ft) and vegetation (aquatic plants)

Comments. For the little minnow, successful spawning depends on having a sustained inundation long enough for the eggs to incubate. These types of scenarios, where any interruption will affect the ecosystem dynamic being modeled, typically use an initial statistical setting of *Minimums* or *Maximums* for the *Duration* query. *Minimums* return a flow that will be that high or higher for the length of the duration. *Maximums* return a flow that will be that low or lower. This relationship also thinks in terms of percentage of years. The *Percent exceedance - Flow frequency* option returns a flow and stage that are equaled or exceeded in the percent of years specified by the user. The *Percent exceedance - Flow duration* option returns a flow and stage that are equaled or exceeded in the percent of time specified by the user.

4.1.2 Big bass winter habitat

A study for big bass showed that fish mortality in a critical over-winter period, **January through May**, was caused by a chronic lack of habitat. Mortalities began to occur when generally poor conditions persisted for more than **two weeks**. Habitat shortages for these fish occur at low flows. Scientists said that these chronic conditions are best represented by **average low flows** and that, since these fish are in the river each winter, using a **typical year (median conditions)** would be a good indicator. The study showed that suitable habitat is proportional to increasing low flows (i.e., higher low flows create more habitat) until those low flows exceed **1,000 cfs**.

HEC-EFM Relationship:

- *Season*: 1/1 to 5/31
- *Duration*: 14 days, *Means* (average) and then *Minimum* (low)
- *Rate of change*: Not applied
- *Percent exceedance*: 50% (2 yr) - *Flow frequency*
- *Hypothesis tracking*: Curve with flow-value points of 0-0, 600-6, 1000-10, 10000-0
- *Geographical queries*: Not applied

Comments. The background for big bass used phrases like "mortality...was caused by a chronic lack of habitat...generally poor conditions". These are clues that big bass are somewhat resilient and will be most affected by bad conditions that occur for long periods of time or sporadically, without enough time between episodes for the fish to recover. This scenario is unlike little minnow spawning where the eggs cannot have bad conditions – be dry – for any single day and survive. Relationships like this typically use an initial statistical setting of *Means* for the *Duration* query.

4.1.3 Benthic macro invertebrate biodiversity

Reservoirs tend to reduce high flows and increase low flows, which creates a more stable flow regime. In these regulated systems, communities of benthic macro invertebrates often have reduced biodiversity because the few species that thrive in the more stable flow conditions out compete all of the others. **Flooding** initiates a return to more natural conditions which encourages the community to rebound to its original biodiversity. Scientists maintain that the **timing is not important**, but the high flows should occur **once every two years**, on average.

HEC-EFM Relationship:

- *Season:* 10/1 to 9/30
- *Duration:* 1 day, *Means* (average) and then *Maximum* (high)
- *Rate of change:* Not applied
- *Percent exceedance:* 50% (2 yr) - *Flow frequency*
- *Hypothesis tracking:* Increased flow will improve (+) benthic biodiversity
- *Geographical queries:* Not applied

Comments. HEC-EFM relationships do not need to be complicated. The background basically says that benthic biodiversity is a function of high flows. So to build this relationship, use statistical queries that focus on high flows of short duration at any time in the water year. Choosing a duration is related to the flashiness of the river system being studied. For the demonstration project, a value of 1 day was chosen, though any duration between 1 and 7 days would likely be a good indicator.

4.1.4 Wetland health

Water exchange between river and wetland areas has also been noted as a key component of wetland health. With frequent exchange, water quality in the wetlands remains good, but with isolation, dissolved oxygen levels drop, wetland areas become anoxic and aquatic species die. This is only an issue in the warm summer months, **mid-May to mid-September**. A hydraulic engineer on your team has determined that flows of **600 cfs** and higher allow water exchange in your project area and a biologist, familiar with the region, suggests that active exchange for approximately **30% of the time** (in summer) will lead to healthy conditions.

HEC-EFM Relationship:

- *Season:* 5/15 to 9/15
- *Duration:* 1 day
- *Rate of change:* Not applied
- *Percent exceedance:* 30% (of time) - *Flow duration*
- *Hypothesis tracking:* Increased flow will improve (+) water exchange for Wetland health
- *Geographical queries:* Not applied

Comments. This relationship is the first demonstration relationship to use the *Percent exceedance - Flow duration* query. In this case, healthy conditions are created when active exchange between the river and wetlands occurs 30% of the time. As mentioned in the little minnow comments, using the *Percent exceedance - Flow duration* option will return a flow and stage that are equaled or exceeded in the percent of time specified by the user.

4.1.5 Riparian tree recruitment and inundation

Reservoir influence and transition of floodplain lands to agriculture has proved a destructive combination for riparian tree forests. Through scientific study, riparian tree establishment has been tied to high flows that occur and recede during germination periods. After germination, survival is a function of water level. If inundated, seedlings are prone to drowning and, conversely, if water levels recede too rapidly, roots desiccate and seedlings are lost.

Germination periods for the riparian tree have been shown to occur between **mid-June through July**. Scientists have found that, after germination, if water levels drop by more than **0.58 feet per week** then riparian tree seedlings will have a lower chance of survival. A high stage needs to occur at least once every **10 years** to keep sustainable riparian tree establishment.

HEC-EFM Relationship: Recruitment

- *Season:* 06/15 to 08/01
- *Duration:* 1 day
- *Rate of change:* 0.58 feet per 7 days - falling (stage)
- *Percent exceedance:* 10% (10 yr) - *Flow frequency*
- *Hypothesis tracking:* Increased flow will improve (+) Riparian tree recruitment
- *Geographical queries:* Not applied

Riparian tree seedlings are sensitive to prolonged periods of inundation. **Beginning of August through mid September** represents the time period immediately following the establishment season, where riparian tree seeds would most likely drown if inundated for an extended length of time. **21 days** is the estimated length of time that a seedling could be continuously inundated before it dies. If the **sustained** inundation is less than 0.5 ft, seedlings are more likely to survive. Median conditions (**2 yr**) can be used as an estimate of typical inundation.

HEC-EFM Relationship: Inundation

- *Season:* 08/01 to 09/15
- *Duration:* 21 days, *Minimums* (sustained highs) and then *Maximum* (extent of effective seedling drowning)
- *Rate of change:* Not applied
- *Percent exceedance:* 50% (2 yr) - *Flow frequency*
- *Hypothesis tracking:* Increased flow will cause Riparian tree inundation to increase (-)
- *Geographical queries:* Depth (0 to 0.5 ft)

Comments. This scenario actually requires two individual relationships that are used to represent a single ecosystem dynamic. The real dynamic of interest is the establishment of new riparian tree seedlings. Establishment occurs when flow and stage conditions are suitable for new seedlings to begin growing (recruit) and when the new seedlings survive inundation that occurs later in the growing season.

The recruitment relationship uses the rate of change query. This query starts at the end of the season and works towards the start of the season one day at a time, checking for violations of the rate of change threshold. The flow and stage at the time of the last violation in the user-defined season is recorded as a “seasonal result”. These seasonal results (one pair of flow and stage for each water year) are then ranked into a frequency table. The statistical result for Riparian tree recruitment is interpolated as the 10% exceedance value.

The inundation relationship uses a *Minimums - Maximum Duration* query to compute the flow and stage that effectively drown or preempt any new recruitment.

These two are used in combination to represent Riparian tree establishment, with the difference between recruitment and inundation results being the portion of recruitment that leads to establishment of new seedlings.

HEC-EFM Combo Relationship: Riparian tree establishment

Primary layer - Riparian tree recruitment

Connected layer - Riparian tree inundation

4.2 Setting up the Model

4.2.1 Creating a New Project and Defining Flow Regimes

1. Open HEC-EFM and use the "File – New" menu option to start a new project. Enter title, author, and project description into the *model information* section of the *Properties Tab*.
2. The dataset for this project is part of the completed demonstration project for HEC-EFM, which was copied to your computer when the HEC-EFM software was installed. Use the "Help – Install Demonstration Project..." menu option to extract the demonstration project to a directory on your computer or network (Figure 10).




Figure 10. Setting up a working directory.

3. Data for the flow regimes are located in "Rolling River Data.dss", which is included in the demonstration project. The first flow regime acts as a placeholder and must be renamed.

Use the “*Edit – Flow Regimes – Rename...*” menu option to open the *Rename Flow Regime* interface. In the *New name* text box, enter Natural and click *OK* (Figure 11).




Figure 11. Naming a flow regime.

4. Add data to this flow regime by moving the cursor over the *Filename* cell and pressing the *browse* button (Figure 12). Browse to the Rolling River folder in the extraction location, select "Rolling River Data.dss", and click *OK*. *Note: The browse button only appears when you have the Filename cell selected.*




Figure 12. Adding the flow regime data file.

- Click the *Open DSS Catalog* button to open the *HEC-DSS Catalog Browser* for that flow regime (Figure 13). Browse to /ROLLING RIVER/AT EFM XS/FLOW//1DAY/NATURAL/, highlight the *f-part* "NATURAL" and click *Select*. HEC-EFM has an auto-mapping feature that looks for stage time series whose *a-*, *b-*, and *f-parts* match the selected flow record. If HEC-EFM finds a match, a message box opens asking whether the matching time series should be used as the corresponding stage values for that flow regime. Click *Yes*. Once both the stage and flow time series have been selected, press *OK*.




Figure 13. Selecting flow and stage time series.

- Use the *Edit – Flow Regimes – Add...* menu option to open the *Add Flow Regime* interface. In the *Name* text box, enter Gaged and click *OK* (Figure 11). Repeat steps 4 and 5 to add data to the new flow regime. Remember to click the *Open DSS Catalog...* button while the cell highlighted in the table of flow regimes is in the row for Gaged and to select the *f-part* "GAGED" after opening the DSS catalog.
- It is important to define the reference flow regime. This is the flow regime that serves as the basis for comparison when looking at direction of ecosystem change. In this example, use the Natural flow regime as the reference. To make a flow regime the reference, click on the box in the *Ref* column of that flow regime (Figure 14). In this case, a check mark is already in the box because Natural was the first flow regime entered. Only active flow regimes are considered during calculations.

Ref	Active	Identifier	Filename	Part A	Part B	Part F	Starting	Ending
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Natural	Rolling River Data..	ROLLING RIVEI	AT EFM X5	NATURAL	01/01/1954	09/30/2003
<input type="checkbox"/>	<input type="checkbox"/>	Gaged	Rolling River Data..	ROLLING RIVEI	AT EFM X5	GAGED	01/01/1954	09/30/2003

Figure 14. Reference and active flow regimes.

- The *Properties Tab* should now look like Figure 15. Use the "File – Save As" menu option to save your project. *Note: The completed demonstration project is called Rolling River.efm. If saving in the extraction directory, please use a different name for your project.*

Figure 15. Completed *Properties Tab* for the demonstration project.

4.2.2 Defining Relationships

- The next step is to enter the relationships. Go to the *Relationships Tab* and use the "Edit – Relationships – Add..." menu option to open the *Add Relationship* interface. In the *Name* text box, enter Little minnow spawning habitat and click *OK* (Figure 16).




Figure 16. Adding a relationship.

2. Enter a description, statistical queries, and hypothesis for the Little minnow spawning habitat relationship (Figure 17).




Figure 17. Statistical queries and hypothesis for the Little minnow relationship.

- Click the *Recalculate* button (bottom right of the interface). Results are displayed on the *Tables Tab* (Figure 18).

Rolling River.efm - HEC-EFM

File Edit Plot Help

Evaluated at: 09/02/2009 14:55

Summary

Relationship	Conf.	Natural	
		Stage, ft	Flow, cfs
Little minnow spawning habitat *		4275.2	1,226

Properties Relationships Tables Combo Relationships

Completed Recalculate

Figure 18. Statistical results for the Natural flow regime and the Little minnow relationship.

- Next, click on the *Properties Tab*; activate the Gaged flow regime by checking its box in the *Active* column of the flow regime table. Click the *Recalculate* button. Results are now displayed for both the Natural and Gaged flow regimes (Figure 19). *Note: It is important to recognize that the Little minnow relationship was not changed during this step. **Flow regimes and relationships exist independently.** When the second flow regime (Gaged) was activated, HEC-EFM computed statistical results for each flow regime using the same statistical criteria, as defined for the Little minnow relationship. Results are different because each flow regime has its own distinct patterns of flow and stage.*

Rolling River.efm - HEC-EFM

File Edit Plot Help

Evaluated at: 09/02/2009 15:01

Summary

Relationship	Conf.	Natural		Chg.	Gaged	
		Stage, ft	Flow, cfs		Stage, ft	Flow, cfs
Little minnow spawning habitat *		4275.2	1,226	Pos	4275.7	1,703

Properties Relationships Tables Combo Relationships

Completed Recalculate

Figure 19. HEC-EFM statistical results of the Little minnow spawning habitat relationship using both the Natural and Gaged flow regimes.

- The next step is to create geographical queries. Go to the *Relationships Tab* and use the "Edit – Manage Geo Queries..." menu option to open the *Manage Tags for Geographical Queries* interface (Figure 20). Click the *Add tag...* button to open the *Add Tag* interface. In the *Name* text box, enter "Depth" and click *OK*. Repeat this step to add another tag named "Vegetation". Now associate the tags with Little minnow spawning habitat by highlighting that relationship in the *Relationship* column and clicking the *Select tags...* button. The *Tag Selector* interface will open. Select *Depth* and *Vegetation* and click the *Add →* button. Click *OK* to save the selections to the *Manage Tags* interface and then click *OK* in the *Manage Tags* interface to save the tags and selections to the project. Text boxes for selected tags will appear on the *Relationships Tab*. Enter a description for each of the Little minnow tags. For *Depth*, enter "Shallow habitat 0 to 3 ft". For *Vegetation*, enter "Requires presence of aquatic plants" (Figure 21).






Figure 20. Adding geographical query tags.

Figure 21. Geographical queries for the Little minnow spawning habitat relationship.



- Repeat steps 1 through 5 for the other relationships (Big bass winter habitat, Benthic macroinvertebrate biodiversity, Wetland health, and Riparian tree recruitment and inundation).
- Turn on *Confidence tracking* for the relationships (Figure 22). For now, leave all relationships at one star.

Figure 22. Confidence tracking and indices.



- Next, create indices by clicking the *Index* button in the *Options* frame (Figure 22). The *Index Description Labels* interface will open. Enter "All" for index A and "Fish" for Index B. Associate each relationship with the appropriate indices by checking box A for each of the relationships and also checking box B for Little minnow and Big bass relationships.
- Your project should now have 2 flow regimes, 6 relationships, and 2 indices. Use the "File – Save" menu option to save your project and then click *Recalculate* to compute results.

4.2.3 Results

Four of the six relationships show a positive change for the Gaged flow regime (Figure 23). Riparian tree inundation and Benthic biodiversity show a negative change. Benthic biodiversity had the most significant change in terms of difference in flow results.

Summary						
Relationship	Conf.	Natural		Chg.	Gaged	
		Stage, ft	Flow, cfs		Stage, ft	Flow, cfs
Little minnow spawning habitat	*	4275.2	1,226	Pos	4275.7	1,703
Big bass winter habitat	*	4274.1	525	Pos	4274.3	609
Benthic macroinvertebrate biodiversity	*	4279.4	6,620	Neg	4277.2	3,190
Wetland health	*	4274.3	636	Pos	4274.5	771
Riparian tree recruitment	*	4274.9	1,017	Pos	4275.1	1,129
Riparian tree inundation	*	4273.7	373	Neg	4274.3	609

Figure 23. Summary of HEC-EFM results.

The index values show a negative response (Neg) for all relationships and a positive response (Pos) for the fish (Figure 24). A positive value suggests that the positive changes outweigh the negatives for the relationships in the index.

Index Values

Index	Gaged
A - All	-4.7
B - Fish	27.4
C -	n/a
D -	n/a
E -	n/a

Figure 24. Index values for “All” relationships and only the relationships that are related to "Fish".

The Rolling River has been a popular study area for wetland research. This has led to numerous peer reviewed studies and an abundance of field data. As these studies support the approach used to define Wetland health, change confidence for that relationship to five stars (Figure 25). Leave all others at 1 star.



Figure 25. Confidence tracking increase for the Wetland health relationship.

Click the *Recalculate* button. Note that the index value for all relationships goes from negative to positive (Figure 26). Indices are best used as screening tools. Lumping multiple, and pseudo-independent, relationships into one numeric index is an approach to be used with caution.


Index Values

Index	Gaged
A - All	9.5
B - Fish	27.4
C -	n/a
D -	n/a
E -	n/a

Figure 26. Index values after a confidence tracking increase for the Wetland health relationship.

Let's also look at Wetland health with a *Reverse lookup* query. To do this, use the "Edit – Relationships – Duplicate..." menu option to open the *Duplicate Relationship* interface. In the *Name* text box, enter Wetland health reverse lookup and click *OK* (Figure 27).

Figure 27. Duplicating a relationship.



Turn on the *Reverse lookup* query by checking its box in the *Other queries* frame. Enter 600 cfs in the text box (Figure 28). Notice that percent exceedance is now grayed out. This is because reverse lookups do not compute a flow based on percent exceedance, instead the user specifies a flow and HEC-EFM computes the percent of years or percent of time that flow is equaled or exceeded. *Note: Non-standard queries (e.g., reverse lookups) cannot be included in the indices.*




Figure 28. Reverse lookup for the Wetland health relationship in HEC-EFM.

Results for Wetland health reverse lookup show that both flow regimes were above the 30% criteria (Figure 29). Results for the Gaged flow regime (67.1%) nearly doubled the 34.1% of time Natural flows in Rolling River equaled or exceeded 600 cfs.

Reverse Look-ups - Flow Duration

Relationship	Conf.	Natural	Gaged	
		% X, of time	Chg.	% X, of time
Wetland health reverse lookup *		34.1	Pos	67.1

Figure 29. Reverse lookup for Wetland health showing percent of time flows above 600 cfs.

4.2.4 HEC-EFM Output

In addition to the statistical results that are output to the *Tables Tab*, HEC-EFM also generates a project summary report and files of the computations it performs while generating the statistical results. To view the project summary report, use the "File – Report" menu option.

By default, computation files are not generated automatically by the software in order to minimize computation time. This output option is selected individually for each relationship. To test this feature, go to the *Relationships Tab*, select the Big bass winter habitat relationship, and check the box next to *Write computation arrays* in the *Options* frame (Figure 30).



Figure 30. Checking the *Write computation arrays* box.

Output formats are selected via the "Edit – Select Arrays Format" menu option. Choices are provided for *XML*, *DSS*, or *Both*. Select *Both* and click the *Recalculate* button (Figure 31). A folder named "Arrays" will be created in the directory of the HEC-EFM project file to store the output.




Figure 31. Output choices via the *Select Arrays Format* menu options.

The term XML stands for Extensible Markup Language. XML is a widely used format for archiving and viewing data. Separate XML files are generated for each combination of flow regime and relationship. Each of these files contains the computational steps, statistical results, and frequency curves produced by HEC-EFM. The XML arrays are organized such that the final results are located at the top of the file followed by each computational step in reverse order (Figure 32). Using the "File – View Array File..." menu option, select "Gaged_Big bass winter habitat.xml" and click *Open*.

The highlighted value is the statistical result. The table also contains seasonal results for each historical water year in the period of record. This type of information provides more details about the relationships and flow regimes than the statistical results reported on the *Tables Tab* and can be very valuable to HEC-EFM applications.

These same data are stored in a file named "arrays.dss" and are archived according to flow regime and relationship names. The "arrays.dss" file can be accessed using HEC-EFM Plotter or HEC-DSSVue, which are available via the HEC website (<http://www.hec.usace.army.mil/>). Figure 33 shows the catalog of data output to DSS for the Big bass winter habitat relationship, time series calculations done to compute seasonal results for the Natural flow regime, and frequency curves of seasonal results for Natural and Gaged flow regimes with the 50% exceedance values selected as the statistical results.

Gaged, Big bass winter habitat

Computations made: 1/22/2008 2:18:30 PM, Rolling River.efm
 Result: Flow = 608.5, Stage = 4,274.273, Closest water year = 1969
 Eco-value = 6.1

Frequency analysis

Plotting position: Weibull

Rank	Frequency	WY	Flow	Stage
1	.0196	1975	868.0	4,274.652
2	.0392	1973	867.9	4,274.688
3	.0588	1984	807.6	4,274.592
4	.0784	1978	776.6	4,274.542
5	.0980	1980	763.6	4,274.522
6	.1177	1977	760.9	4,274.518
7	.1373	1976	745.4	4,274.493
8	.1569	1983	742.4	4,274.488
9	.1765	1998	735.9	4,274.478
10	.1961	1972	705.9	4,274.430
11	.2157	1990	693.3	4,274.406
12	.2353	1964	692.1	4,274.407
13	.2549	1965	683.9	4,274.394
14	.2745	1966	675.1	4,274.380
15	.2941	1958	668.4	4,274.370
16	.3137	1991	666.5	4,274.365
17	.3333	1954	657.3	4,274.352
18	.3529	1979	650.8	4,274.341
19	.3726	1993	650.1	4,274.333
20	.3922	1960	642.6	4,274.328
21	.4118	1974	638.6	4,274.322
22	.4314	1970	637.6	4,274.320
23	.4510	1971	632.2	4,274.312
24	.4706	1968	624.3	4,274.299
25	.4902	1969	612.6	4,274.280
25	.5000		608.5	4,274.273
26	.5098	1967	604.5	4,274.267
27	.5294	1982	588.8	4,274.242

This is the statistical result for the Big bass winter habitat relationship and Gaged flow regime. A statistical result is the pair of flow and stage that meet the statistical criteria used to define the relationship.

Figure 32. Excerpt from the XML output file for Big bass winter habitat and the Gaged flow regime.




Figure 33. Catalog of HEC-EFM data output to DSS for the Big bass winter habitat relationship. Seasonal results for the Natural flow regime as well as a frequency curve of seasonal results for Natural and Gaged flow regimes are shown.

4.2.5 Using HEC-EFM Plotter

HEC-EFM Plotter is designed to help users view output and compare results for different flow regimes and relationships. Additionally, by displaying each computational step that HEC-EFM performs while analyzing time series, HEC-EFM Plotter offers an opportunity to understand the statistical process and settings being used to investigate each relationship.

Initiate HEC-EFM Plotter by selecting the “Plot – Activate HEC-EFM Plotter” menu option. DSS output is automatically imported to *Standard Plots* for each combination of flow regime and relationship. Use the *Relationship* dropdown list in the upper left hand corner of the main interface to select the Big bass winter habitat relationship (Figure 34). Choose which flow regime (e.g. Natural or Gaged) to view using the *Flow Regime* dropdown list.




Figure 34. HEC-EFM Plotter showing results for the Big bass winter habitat relationship with the Natural flow regime.

Custom Plots are useful when comparing results for multiple relationships or flow regimes. Select the Custom Plots option and add a Custom Plot called “Bass Compare” by choosing Add... from the Custom Plots dropdown button (Figure 35).




Figure 35. Adding a custom plot.

Click the DSS Selector button and browse to the arrays.dss file for your project. Select all records with a b-part of Big bass winter habitat and click the Set Pathname button (Figure 36) to import those data to the “Bass compare” plot (Figure 37).




Figure 36. Process for selecting Big bass winter habitat results for comparison in a custom plot within HEC-EFM Plotter.




Figure 37. Custom plot in HEC-EFM Plotter comparing the Big bass winter habitat relationship for Gaged and Natural flow regimes.

HEC-EFM Plotter has many features for navigating (zoom, hover, snap to water year, advance/retreat a year) and viewing data sets (plot markers, show/hide/reorder data sets via the legend, axis settings, toggle plots with a Ctrl-T command).

HEC-EFM Plotter and HEC-EFM can be open at the same time. In fact, when a setting is changed in HEC-EFM and a *Recalculate* is performed, plots can be updated by clicking the *Refresh* button, which makes HEC-EFM Plotter an effective way for teams to explore and refine the statistical settings that define the relationships.

4.3 Hydraulic Analysis

Spatial analysis of HEC-EFM results can be done using water surface profiles and grids of depth and velocity produced by a geo-referenced hydraulic model. Past applications of HEC-EFM have used HEC-RAS and HEC-GeoRAS, which can be found on HEC's website <http://www.hec.usace.army.mil/>. Documentation on the use of HEC-RAS and HEC-GeoRAS are also available at the website.

4.3.2 HEC-GeoRAS

HEC-GeoRAS is a set of procedures, tools, and utilities for pre and post-processing geospatial data in ArcGIS® (Figure 39). With HEC-GeoRAS, water surface profile data exported to GIS can be used with a digital terrain model to calculate depth grids, velocity grids, and floodplain boundary polygons.



Figure 39. Example of a depth grid computed in HEC-GeoRAS using the water surface profile simulated by HEC-RAS for the statistical results of Riparian tree inundation from HEC-EFM.

4.4 HEC-EFM Analyses with GIS

Geographic information systems are technologies used to store, manage, edit, analyze, and display data that are spatially referenced to the earth. Using GIS in an application of HEC-EFM allows users to analyze data layers produced by HEC-GeoRAS as well as any external data sets that have ecological significance. The following sequence of figures is an example of how GIS can be used to investigate ecosystem response spatially. Results are shown for Riparian tree establishment dynamics in the Natural flow regime. Depth grids were initially created for recruitment (Figure 40) and inundation (Figure 41). The inundation layer was then placed on top of the recruitment layer (Figure 42).



Figure 40. Depth grid created for Riparian tree recruitment. The green area shows where stage recession creates suitable conditions for seedlings to begin to grow.



Figure 41. Depth grid created for Riparian tree inundation. This blue area shows where any new seedlings will be drowned by prolonged inundation.




Figure 42. Overlay of Riparian tree recruitment and Riparian tree inundation depth grids.

In accordance with the combo relationship for Riparian tree establishment (Figure 6), the inundation depth grid was clipped from the recruitment depth grid. The resulting layer shows the area where HEC-EFM predicts Riparian tree establishment (Figure 43).



Figure 43. Depth grid created by clipping the inundation depth grid from the recruitment depth grid. The resulting area shows where HEC-EFM predicts Riparian tree establishment.

Information for the Riparian tree inundation relationship mentioned that seedlings were more likely to survive if sustained inundation did not exceed 0.5 ft. Delineating the area of inundation less than or equal to 0.5 ft shows the portion of Riparian tree inundation that may not lead to seedling mortality (Figure 44). This Riparian tree inundation fringe can then be layered with the clipped layer to create a new view of predicted and possible (due to the shallow depths) Riparian tree establishment (Figure 45).




Figure 44. The yellow fringe shows areas of the Riparian tree inundation depth grid that are less than or equal to 0.5 ft, which may not be deep enough to drown new seedlings.




Figure 45. Potential recruitment sites for the Rolling River's Natural flow regime as computed using statistical and geographical queries in HEC-EFM, water surface profiles computed by HEC-RAS, and depth grids produced by HEC-GeoRAS.

So far the demonstration project has focused on comparing flow regimes with different hydrologic scenarios (Natural vs. Gaged). HEC-EFM can also be used to look at changes in topography, which will change the stage time series without changing the flow. Figure 46 shows a comparison of the current topography and a planned restoration project that increases stream meander. The same flows (Natural) are used for each of the channel topographies.




Figure 46. Comparison of Riparian tree establishment in existing and restored topographic conditions.

The following progression of tasks for the demonstration project has now been completed: 1) building relationships, 2) setting up an HEC-EFM project and computing statistical results, 3) simulating the statistical results with a hydraulic model, and 4) use of GIS to investigate ecosystem responses spatially.

Statistical results in HEC-EFM are computed as flows and stages, percent exceedances, index values, and directions of ecosystem change. Spatial results are shown as combinations of map layers with each combination presenting a snapshot of an ecosystem dynamic. Layers are typically translated to areas for spatial comparisons of different flow regimes and channel topographies. Working with GIS allows use of other ecologically relevant data. For example, a soils map for Rolling River could be used to intersect the Riparian tree establishment layer (Figure 35) with the extent of suitable soils for riparian trees.

CHAPTER 5

Conclusions

This quick start guide is designed to demonstrate the process used to apply HEC-EFM and to highlight some features of the software. Everything covered in this document and all features active in the software have been tested extensively, but there is still the potential for undetected problems to surface. If a bug or suspect behavior is noted, please refer to HEC's website (www.hec.usace.army.mil/software/HEC-EFM/BugReporting/) for guidance on how to report bugs.

HEC-EFM is still a relatively young software tool. Several new features are being actively developed. Two exciting and ongoing efforts are HEC-GeoEFM and the development of a continuous simulation capability for HEC-EFM.

HEC-GeoEFM is a partnered effort between HEC and Environmental Systems Research Institute, Inc. (ESRI) to develop a tool that manages the spatial data and results of HEC-EFM projects. It will also contain a package of features to help users perform the spatial calculations typically done in HEC-EFM studies.

The continuous simulation feature is a long-term development direction that will ultimately perform simulations of ecosystems in time and space (animations of ecosystems).

Look for these and other new features in future versions of HEC-EFM.

CHAPTER 6

Terms and Conditions for Use

Use of the program is governed by the terms and conditions for use. They limit what can be done with the program software, waive warranty, limit liability, and indemnify the developers and the United States government. The program cannot be used unless the terms and conditions for use are accepted; the full text is given below.

Terms and Conditions for Use

The United States Government, US Army Corps of Engineers, Hydrologic Engineering Center ("HEC") grants to the user the rights to install Ecosystem Functions Model (HEC-EFM) "the Software" (either from a disk copy obtained from HEC, a distributor or another user or by downloading it from a network) and to use, copy and/or distribute copies of the Software to other users, subject to the following Terms and Conditions for Use:

All copies of the Software received or reproduced by or for user pursuant to the authority of this Terms and Conditions for Use will be and remain the property of HEC.

User may reproduce and distribute the Software provided that the recipient agrees to the Terms and Conditions for Use noted herein.

HEC is solely responsible for the content of the Software. The Software may not be modified, abridged, decompiled, disassembled, un-obfuscated or reverse engineered. The user is solely responsible for the content, interactions, and effects of any and all amendments, if present, whether they be extension modules, language resource bundles, scripts or any other amendment.

The name "HEC-EFM" must not be used to endorse or promote products derived from the Software. Products derived from the Software may not be called "HEC-EFM" nor may any part of the "HEC-EFM" name appear within the name of derived products.

No part of this Terms and Conditions for Use may be modified, deleted or obliterated from the Software.

No part of the Software may be exported or re-exported in contravention of U.S. export laws or regulations.

Waiver of Warranty

THE UNITED STATES GOVERNMENT AND ITS AGENCIES, OFFICIALS, REPRESENTATIVES, AND EMPLOYEES, INCLUDING ITS CONTRACTORS AND SUPPLIERS PROVIDE HEC-EFM `"AS IS,"` WITHOUT ANY WARRANTY OR CONDITION, EXPRESS, IMPLIED OR STATUTORY, AND SPECIFICALLY DISCLAIM ANY IMPLIED WARRANTIES OF TITLE, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE AND NON-INFRINGEMENT.

Depending on state law, the foregoing disclaimer may not apply to you, and you may also have other legal rights that vary from state to state.

Limitation of Liability

IN NO EVENT SHALL THE UNITED STATES GOVERNMENT AND ITS AGENCIES, OFFICIALS, REPRESENTATIVES, AND EMPLOYEES, INCLUDING ITS CONTRACTORS AND SUPPLIERS, BE LIABLE FOR LOST PROFITS OR ANY SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES ARISING OUT OF OR IN CONNECTION WITH USE OF HEC-EFM REGARDLESS OF CAUSE, INCLUDING NEGLIGENCE.

THE UNITED STATES GOVERNMENT'S LIABILITY, AND THE LIABILITY OF ITS AGENCIES, OFFICIALS, REPRESENTATIVES, AND EMPLOYEES, INCLUDING ITS CONTRACTORS AND SUPPLIERS, TO YOU OR ANY THIRD PARTIES IN ANY CIRCUMSTANCE IS LIMITED TO THE REPLACEMENT OF CERTIFIED COPIES OF HEC-EFM WITH IDENTIFIED ERRORS CORRECTED. Depending on state law, the above limitation or exclusion may not apply to you.

Indemnity

As a voluntary user of HEC-EFM you agree to indemnify and hold the United States Government, and its agencies, officials, representatives, and employees, including its contractors and suppliers, harmless from any claim or demand, including reasonable attorneys' fees, made by any third party due to or arising out of your use of HEC-EFM or breach of this Agreement or your violation of any law or the rights of a third party.

Assent

By using this program you voluntarily accept these terms and conditions. If you do not agree to these terms and conditions, uninstall the program and return any program materials to HEC (If you downloaded the program and do not have disk media, please delete all copies, and cease using the program.)